FA: Quoi faire quand on ne peut plus anticoaguler?

Montréal, 21 septembre 2018

Dr Blandine Mondésert
Electrophysiology service, Medicine department
Adult Congenital Heart Disease Center
Montreal Heart Institute, Université de Montréal

Conflits d'intérêt

- Bourses de recherche:
 - ✓ Boston Scientific
- Honoraires/consultants:
 - ✓ Boston Scientific
 - ✓ Medtronic
 - ✓ Abbott
 - ✓ Servier
 - ✓ Boehringer
 - ✓ Bayer

Objectifs

- Connaître les options prévenant l'AVC pour la FA pour le patient non « anticoagulable »
- Savoir sélectionner le patient pour les approches invasives et comment référer en spécialité
- Comprendre l'essentiel de la technique et le suivi requis dans la communauté.

Cas Clinique n1

- Homme de 76 ans
- FA non valvulaire
- Diabète, HTA contrôlée, coronarien
- Fonction rénale normale (GFR > 60)

Cas Clinique n1

Case #1 – Sequence of events

Saignements digestifs

Apixaban = no more GI bleeding than warfarin (12% less, p=NS)

Dabigatran 150 BID and rivaroxaban 20 mg die = at least 40% more GI bleeding than warfarin (p=signif)

Edoxaban 60 mg die = + 20% (p=NS)

Cas Clinique n1

Recommanderiez-vous une fermeture d'appendice auriculaire?

A. Oui

B. Non

Cas Clinique n2

- Homme de 72 ans
- HTA
- FA non valvulaire
- GFR > 60

Cas Clinique n2

Case #2 – Sequence of events

Cas Clinique n2

- A. Je ré-anticoagule le patient (patient réfractaire)

- B. J'arrête l'anticoagulation sans fermer l'appendice auriculaire

- C. J'arrête l'anticoagulation et réfère la patient pour fermeture de l'appendice auriculaire

Cas Clinique n2

Case #2 – Sequence of events

Thrombus

Localisation du Thrombus dans la FA non-valvulaire:
91% (201/222) LAA

Setting	N	Appendage	%	LA Body	%	Reference
TEE	317	66	21	1	0.3	Stoddard; JACC, 1995
TEE	233	34	15	1	0.4	Manning; Circ, 1994
Autopsy	506	35	7	12	2.4	Aberg; Acta Med Scan, 1969
TEE	52	2	4	2	3.8	Tsai; JFMA, 1990
TEE	48	12	25	1	2.1	Klein; Int J Card Image, 1993
TEE & Operation	171	8	5	3	1.8	Manning; Circ, 1994
SPAF III TEE	359	19	5	1	0.3	Klein; Circ, 1994
TEE	272	19	7	0	0.0	Leung; JACC, 1994
TEE	60	6	10	0	0.0	Hart; Stroke, 1994
Total Thrombus		201		21

Thrombus

Le caillot qui se forme dans l'auricule est souvent volumineux = AVC significatif

Migration d'un thrombus de l'AAG

Cas Clinique n3

- Femme de 77 ans
- STEMI avec DES sur IVA
- Diabète, tabagisme actif
- FA sous Rivaroxaban 20mg
- Fonction rénale normale (GFR>60)

Cas Clinique n3

Quel traitement?

- A. ASA+CLOPIDOGREL+OAC
- B. CLOPIDOGREL+OAC
- C. Fermeture AAG pour permettre ASA+TICAGRELOR

Cas Clinique n4

- Femme de 83 ans
- Référée pour TAVI
- FA permanente sous Apixaban 2.5mg BID
- Coronarienne, diabète, hypertension
- Insuffisance rénale chronique

Cas Clinique n4

Cas Clinique n4

Quelle est l'incidence d'un thrombus dans l'AAGau CT scan chez les patients référés pour TAVI?

- A. <5%
- B. >10%
- C. Pas de relation avec des VAC périprocéduraux

Cas Clinique n4

Quelle est l'incidence d'un thrombus dans l'AAGau CT scan chez les patients référés pour TAVI?

- A. <5%
- B. >10%: 11% chez tous les patients, 32% chez les patients en FA
- C. Pas de relation avec des VAC périprocéduraux

Limitations des DOACs

- Risque intrinsèque de saignement
- Compliance
- Coût
- Prescription limitée par la fonction rénale
- Pas d'antidote sauf pour le dabigatran...

Limitations des DOACs

- Risque intrinsèque de saignement
- Compliance
- Coût
- Prescription limitée par la fonction rénale
- ~~Pas d'antidote sauf pour le dabigatran...~~

Impact du saignement

- Symptômes chroniques: fatigue, anémie, insuffisance cardiaque
- Hospitalisations
- Transfusions
- Coût et risques des tests

Les patients les plus à risque

- ATCD d'hémorragie cérébrale
- Angiopathie amyloïde cérébrale, micro-saignements cérébraux
- Cirrhose, saignements digestifs récurrents (angiodysplasie, source inconnue, post radiation)
- Insuffisance rénale sous dialyse, hématurie récidivante (tumeur)
- Rendu-Osler-Weber

Procédure fermeture d'AG

Développée comme **alternative à l'ACO** chez les patients avec

- **FA non valvulaire** à risque d'événements thrombo-emboliques (CHADS ≥ 1)
- **qui ne tolèrent pas l'anticoagulation ou qui sont à haut risque de saignement**

Options pour exclusion d'auricule

- Chirurgie
- Approches percutannées
 - ✓ Méthodes endocavitaires avec installation de prothèses
 - ✓ Méthodes extra-cardiaques avec installation de suture (LARIAT)

Différentes formes d'AAG

Différentes formes d'AAG

The Wind Sock Type LAA is an anatomy in which one dominant lobe of sufficient length is the primary structure.

The Chicken Wing Type LAA is an anatomy whose main feature is a sharp bend in the dominant lobe of the LAA anatomy at some distance from the perceived LAA ostium.

The Broccoli Type LAA is an anatomy whose main feature is an LAA that has limited overall length with more complex internal characteristics.

Prothèses intra-cardiaques

- Watchman (Atritech $\square$ Boston Scientific)
- AMPLATZER Cardiac Plug - Amulet (2^e génération (AGA $\square$ St-Jude Medical $\square$ Abbott)

CE mark since 2005
FDA approved since March 2015

CE mark since 2008
FDA approved since 2017

Même but, philosophie différente

AMPLATZER Amulet
(Abbott)

"pacifier principle"¹

Watchman
(Boston Scientific)

"half football principle"²

Park at CSI 2011. "Current Status of Left Atrial Appendage Closure Devices – Amplatzer ACP Occluder", Frankfurt, June 23-25, 2011.

Bénéfices

- Chez des patients **NON ELIGIBLE** à l'anticoagulation
 - ✓ Réduction de 80% du risque d'AVC hémorragique à 5 ans
 - ✓ Etudes randomisées WATCHMAN contre warfarine: PROTECT-AF et PREVAIL

Bénéfices

Long-Term (5 years) Patient-Level Meta-Analysis of PROTECT AF and PREVAIL

Figure 4 Actual major bleeding rate excluding procedural bleeding events, and calculated major bleeding risk based on HAS-BLED score, and the relative risk reduction for the total EWOLUTION patient cohort after 1-year follow-up. RR = relative risk; SE = systemic embolism; TIA = transient ischemic attack.

Bénéfices

- Chez des patients NON ELIGIBLE à l'anticoagulation
 - ✓ Réduction de 60% du risque attendu d'AVC et d'embolies systémiques
 - ✓ Registres: ASAP et EWOLUTION, ACP et AMULET

Comparaison des registres

	ACP Registry ¹	Watchman EWOLUTION ²	Amulet (Current Study)
Implant Success	97.3%	98.5%	98.8%
LAA Closure Rate (1-3 months) $\leq$ 5 mm	98.1%	99.3%	100.0%
Device or Procedure- Related Complications	5.0%	2.7%	2.7%
Early Mortality	0.8% (30-day)	0.7% (30- day)	0.3% (7-day)

¹ Tzikas et al. *EuroIntervention*. 2015;10

² Boersma et al. *Eur Heart J*. 2016 Aug;37(31):2465-74

Figure 3 Actual stroke rate, and calculated stroke risk based on CHA₂DS₂-VASc score, and the relative risk reduction for the total EWOLUTION patient cohort after 1-year follow-up. RR = relative risk; SE = systemic embolism; TIA = transient ischemic attack.

Expérience Canadienne

Table 8. Follow-up cardiovascular events

Overall Events	N=106
Mean follow-up (days)	210.3 ± 182.2
- CV death	2 (1.9%)
- Non-CV death	2 (1.9%)
- Stroke	0 (0%)
- TIA	2 (1.9%)
- Systemic embolization	0 (0%)
- MI	1 (1.0%)
- Major bleeding	5 (4.7%)*
- Device associated thrombus	1 (1.0%)

*3 on DAPT, 1 on warfarin, 1 on ASA alone

Table 3 Listing of all major cardiac adverse events within 7 d of implant and other device-/procedure-related serious adverse events

Major adverse cardiac events* ≤ 7 d	18 (1.8%)
All deaths (0-7 d) [†]	4
Major bleeding	9
Cardiac tamponade/significant PE	3
Device embolization requiring surgery	1
Device embolization snared	1
Stroke	None
Systemic embolism	None
Myocardial infarction	None
Other events requiring surgery/major intervention	None
Other periprocedural serious adverse events ≤ 7 d	15 (1.5%)
Vascular complications @ groin	4
Air embolism (coronary)	2
Minor pericardial effusion (untreated)	2
Reinterventions due to incomplete seal	2
Minor bleeding (untreated)/hematoma	2
TIA	1
Hypotension	1
Adverse reaction to anesthesia	1

PE = pericardial effusion; TIA = transient ischemic attack.

*Major cardiac adverse events include 5 events (3 deaths and 2 major bleeding events) that are deemed unrelated to the procedure, yet we include them in the list because of their proximity to the implant procedure.

[†]One event >7 d but originating at the procedure.

Expérience Canadienne

Table 7. Major procedural safety events

Major Procedural Safety Events	N=106
Major in-hospital or within 7days	
- Death*	1 (0.9%)
- Stroke/TIA/Systemic Embolization	0 (0%)
- MI	0 (0%)
- Cardiac tamponade*	1 (0.9%)
- Device embolization (snared)	1 (0.9%)
<i>Composite major safety events</i>	<i>2 (1.9%)</i>
Other in-hospital events	
- Pericardial effusion (small, no drainage)	2 (1.9%)
- Asymptomatic Hg drop (transfusions)	2 (1.9%)
- Minor bleed (hematoma)	1 (0.9%)
- Pseudoaneurysm (thrombin injection)	1 (0.9%)

*required surgical repair, died 5 days later from sepsis

ESC Guidelines FA

Occlusion or exclusion of the left atrial appendage

Recommendations	Class	Level
After surgical occlusion or exclusion of the LAA, it is recommended to continue anticoagulation in at-risk patients with AF for stroke prevention.	I	B
LAA occlusion may be considered for stroke prevention in patients with AF and contra-indications for long-term anticoagulant treatment (e.g. those with a previous life-threatening bleed without a reversible cause).	IIb	B
Surgical occlusion or exclusion of the LAA may be considered for stroke prevention in patients with AF undergoing cardiac surgery.	IIb	B
Surgical occlusion or exclusion of the LAA may be considered for stroke prevention in patients undergoing thoracoscopic AF surgery.	IIb	B

European Heart Journal (2016) 37, 2893–2962 doi:10.1093/eurheartj/ehw210

www.escardio.org/guidelines

European Heart Journal - doi:10.1093/eurheartj/ehw210

EUROPEAN
SOCIETY OF
CARDIOLOGY®

INSTITUT DE
CARDIOLOGIE
DE MONTRÉAL

Etudes à venir

ASAP-TOO (NCT02928497): Overview

WATCHMAN™
LEFT ATRIAL APPENDAGE
CLOSURE DEVICE

Study Objective	Evaluate LAA Closure with WATCHMAN in NVAF patients deemed not suitable for oral anti-coagulation therapy
Study Design	Prospective, multi-center Randomized 2:1 (Watchman vs Control) Considering Group Sequential Design
Primary Endpoint	<p><u>Effectiveness Endpoint</u> Time to first occurrence of ischemic stroke or systemic embolism</p> <p><u>Safety Endpoint</u> 7-day rate of all-cause death, ischemic stroke, systemic embolism, or device- or procedure- related events requiring open cardiac surgery or major endovascular intervention</p>
Patient Population	888
Number of Sites	100 global sites
Follow-up*	<ul style="list-style-type: none"> 45 Day with TEE 6,18 month phone visit 12 month with TEE Bi-annually for years 2-5

* Brain imaging required at baseline if prior stroke or TIA

AMULET IDE TRIAL

Brief Summary:

The Amulet™ device will be evaluated for safety and efficacy by demonstrating its performance is non-inferior to the commercially available WATCHMAN® left atrial appendage closure device in patients with non-valvular atrial fibrillation. Patients who are eligible for the trial will be randomized to receive either the Amulet device or the WATCHMAN device and will be followed for 5 years after device implant.

Condition or disease ⓘ	Intervention/treatment ⓘ	Phase ⓘ
Stroke	Device: Amulet Left Atrial Appendage Occluder Device: WATCHMAN Left Atrial Appendage Closure	Not Applicable

Detailed Description:

The Amulet IDE trial is a prospective, randomized, multi-center active control worldwide trial, designed to evaluate the safety and effectiveness of the AMPLATZER™ Amulet™ Left Atrial Appendage Occluder. Subjects will be randomized in a 1:1 ratio between the Amulet LAA occlusion device (treatment) or a Boston Scientific WATCHMAN® LAA closure device (Control). The trial will be conducted at up to 150 sites worldwide. All enrolled subjects will follow the protocol-required tests and assessments at each scheduled follow-up visit.

Study Design

Go to

Study Type ⓘ: Interventional (Clinical Trial)

Estimated Enrollment ⓘ: 1878 participants

Allocation: Randomized

Intervention Model: Parallel Assignment

Masking: None (Open Label)

Primary Purpose: Prevention

Official Title: AMPLATZER™ Amulet™ Left Atrial Appendage Occluder Randomized Controlled Trial

Study Start Date ⓘ: August 2016

Estimated Primary Completion Date ⓘ: February 2020

Estimated Study Completion Date ⓘ: December 2023

NCT02879448

INSTITUT DE
CARDIOLOGIE
DE MONTRÉAL

Etudes à venir

LAOOS III Trial NCT01561651

(PI = Richard Whitlock, McMaster University)

- Surgical LAAC with suture and/or stapler or a regulatory approved device vs standard therapy
- Inclusion:
 - 1. Greater than 18 years of age
 - 2. Undergoing cardiac surgical procedures with the use of cardiopulmonary bypass
 - 3. Have a documented history of atrial fibrillation or atrial flutter
 - 4. CHA2DS2-VASc score ≥ 2
- Primary outcomes:
 - Stroke or systemic arterial embolism [Time Frame: Common termination point (median follow-up of 4 years)]
 - First occurrence of ischemic stroke or transient ischemic attack with positive neuroimaging or systemic arterial embolism

En pratique

- Le plus souvent chez patients dont l'anticoagulation a été cessée (pas de liste d'attente)
- ETO et/ou CT scanner quelques jours avant pour r/o un thrombus et définir l'anatomie
- Hospitalisation de courte durée

En pratique

- Sous anesthésie générale et guidance ETO
- Alternative = Anesthésie locale et ICE
- Ponction trans-septale – 40 cc de colorant

Procédure

Implant procedure performed
with both Fluoroscopy and TEE

Content of this animation is for information purposes only and not meant for medical diagnostic. Information for use only in countries with applicable health authority product registrations.
Information contained herein for distribution outside the USA, France and Japan only. © 2014 Boston Scientific Corporation or its affiliates. All rights reserved.

Post-procédure

- ETO à 4-6 semaines pour exclure thrombus et/ou fuite résiduelle
- Prophylaxie de l'endocardite infectieuse
1 an

Thrombus post-procédure

TABLE 2 Major Adverse Events (n = 98) in Patients Treated With LAA Occlusion Using the Nitinol Plug or Nitinol Cage Devices

	Overall (N = 469)	Nitinol Cage (n = 272)	Nitinol Plug (n = 197)	p Value (Nitinol Cage vs. Nitinol Plug)
Death	33 (6.9)	18 (6.7)	15 (7.1)	0.85
Ischemic stroke	19 (4.0)	10 (3.7)	9 (4.3)	0.86
TIA	2 (0.4)	2 (0.7)	0 (0)	—
Major hemorrhage	18 (3.8)	10 (3.7)	8 (3.8)	0.76
Thrombus on the device				
In the whole study group	26 (5.4)	13 (4.8)	13 (6.2)	0.36
In patients with LAA imaging	26 (7.2)	13 (5.5)	13 (11.0)	0.02

Values are n (yearly rate %).

LAA – left atrial appendage; TIA – transient ischemic attack.

TABLE 4 Multivariable Analysis (Cox Regression Model) for Predictors of Thrombus Formation on the Device and Predictors of Stroke and TIA*

	HR (95% CI)	p Value
Thrombus formation on the device		
Age (per 1-yr increase)	1.07 (1.01-1.14)	0.02
Previous ischemic stroke	3.68 (1.17-11.62)	0.03
CHA ₂ DS ₂ -VASc score	0.69 (0.44-1.06)	0.09
APT at discharge	0.35 (0.12-1.04)	0.06
Dual APT at discharge	0.10 (0.01-0.76)	0.03
OAC at discharge	0.26 (0.09-0.77)	0.02
Strokes or TIAs		
Vascular disease	5.03 (1.39-18.23)	0.01
Thrombus on the device	4.39 (1.05-18.43)	0.04
CHA ₂ DS ₂ -VASc score	0.71 (0.47-1.06)	0.09
APT at discharge	1.35 (0.20-9.06)	0.75
Dual APT at discharge	0.64 (0.15-2.69)	0.54
OAC at discharge	0.39 (0.06-2.61)	0.33

*Analysis restricted to patients with LAA imaging during follow-up. For prediction of thrombus formation on the device, time zero is time at discharge after LAA closure. For prediction of stroke or TIA, time zero is time at first post-procedure LAA imaging.

CI – confidence interval; HR – hazard ratio; other abbreviations as in [Table 1](#).

Thrombus post-procédure

CENTRAL ILLUSTRATION Kaplan-Meier Cumulative Event-Free Curves of Ischemic Strokes and Transient Ischemic Attacks With and Without Thrombus on the Device

Ischemic Stroke / Transient Ischemic Attack

N at risk	0	6	12	18	24
No thrombus	300	100	72	62	39
Thrombus	25	12	6	2	0

Fauchier, L. et al. *J Am Coll Cardiol.* 2018;71(14):1528-36.

The curves are representative of being event-free for ischemic strokes and transient ischemic attacks, with and without thrombus on the device, after left atrial appendage occlusion. Time zero is time at first post-procedure left atrial appendage imaging. The curves demonstrate a higher risk for ischemic strokes or transient ischemic attacks in the patients with a diagnosis of device-associated thrombus after left atrial appendage occlusion. The mean follow-up time was 13 ± 13 months. CI = confidence interval; HR = hazard ratio.

FIGURE 1 Flowchart of the Study Group

The study cohort from RELEXAO (Registry on Real-Life Experience With Left Atrial Appendage Occlusion: [NCT03279406](https://clinicaltrials.gov/ct2/show/study/NCT03279406)) between February 1, 2007 and January 31, 2017 is shown. The proportions of the different devices for left atrial appendage (LAA) closure and patients with left atrial appendage imaging are shown, as well as the different antithrombotic regimens at hospital discharge. AF = atrial fibrillation; APT = antiplatelet therapy; OAC = oral anticoagulation.

Expérience Canadienne

Discharge Antithrombotic Agents	N=103
Aspirin	87 (84.5%)
Clopidogrel	80 (77.7%)
Ticagrelor	1 (1.0%)
Warfarin	6 (5.8%)
Direct OAC	16 (15.5%)
Heparin	0 (0%)
DAPT alone	76 (73.4%)*
Aspirin alone	3 (2.9%)
Warfarin alone	2 (1.9%)
Warfarin + (Aspirin or Clopidogrel)	3 (2.9%)
Warfarin + DAPT	1 (1.0%)
Direct OAC alone	12 (11.7%)
Direct OAC + Aspirin	1 (1.0%)
Direct OAC + DAPT	3 (2.9%)
None	2 (1.9%)

*ASA+clopidogrel, except 1 on ASA+ticagrelor

Antithrombotic Agents Duration	N=103
DAPT	76 (73.8%)
- 1mth	6 (5.8%)
- 3mth	57 (55.3%)
- 6mth	14 (13.6%)
- >6mth	3 (2.9%)
Warfarin or DOAC	21 (20.4%)
- 6weeks	12 (11.7%)
- 3mth	1 (1.0%)
- 6mth	3 (2.9%)
- >6mth	2 (1.9%)

Thrombus post-procédure

Table 3. Event Rates for Patients With or Without Device-Related Thrombus

Event	Device-Related Thrombus		Unadjusted		Adjusted	
	No (n=1674)	Yes (n=65)	Rate Ratio (95% CI)	P Value	Rate Ratio (95% CI)	P Value
All stroke or systemic embolism >0 days postimplant	1.78 (123/6904.8)	7.46 (19/254.7)	4.15 (2.55–6.75)	<0.001	3.55 (2.18–5.79)	<0.001
Ischemic stroke or systemic embolism >0 days postimplant	1.65 (114/6904.8)	6.28 (16/254.7)	3.75 (2.22–6.35)	<0.001	3.22 (1.90–5.45)	<0.001
Ischemic stroke >0 days postimplant	1.65 (114/6904.8)	5.10 (13/254.7)	3.05 (1.71–5.43)	<0.001	2.60 (1.46–4.64)	0.001
Hemorrhagic stroke >0 days postimplant	0.13 (9/6904.8)	1.18 (3/254.7)	9.04 (2.44–33.40)	0.001	7.98 (2.12–29.94)	0.002
Disabling or fatal stroke >0 days postimplant	0.31 (21/6847.5)	1.22 (3/246.0)	3.78 (1.12–12.72)	0.032	3.14 (0.93–10.62)	0.066
Cardiovascular/unknown cause death	1.78 (123/6904.8)	2.75 (7/254.7)	1.53 (0.71–3.28)	0.278	1.31 (0.61–2.82)	0.486
All-cause death	4.43 (306/6904.8)	5.89 (15/254.7)	1.34 (0.80–2.25)	0.272	1.13 (0.67–1.90)	0.649
Major bleeds	4.11 (284/6904.8)	7.85 (20/254.7)	1.95 (1.24–3.08)	0.004	1.78 (1.13–2.81)	0.013

All results come from a generalized linear mixed model with a random study effect. Adjusted results are adjusted for CHA₂DS₂-VASc and HAS-BLED scores. CI indicates confidence interval.

Thrombus post-procédure

Figure 3. Kaplan-Meier survival curves for thromboembolic events as a function of device-related thrombus (DRT).

A. The presence of DRT is associated with higher rates of ischemic stroke or systemic embolism. **B.** Similarly, the rates of all-cause stroke or systemic embolism are also higher if DRT is present. CI indicates confidence interval; and HR, hazard ratio.

Thrombus post-procédure

Figure 6. Ischemic stroke rates after Watchman implant versus expected rate with oral anticoagulants.

Anticoagulants/Anti-agrégants post procédure?

- Si patient était anticoagulé:
 - ✓ ASA à vie
 - ✓ ACO 6 semaines (Watchman)

- Si patient n'était plus anticoagulé:
 - ✓ Aspirine à vie si possible
 - ✓ Clopidogrel 6-12 semaines

- La vraie vie....

Conclusions

- Thérapie locale alternative raisonnable à l'anticoagulation orale (un traitement systémique)
- Prévenir les AVCs chez les pts avec FA à haut risque thrombo-embolique **ET** à haut risque de saignement
- Avant d'envisager un remplacement diffus de l'anticoagulation en pratique clinique
- Comparaison de l'efficacité des prothèses vs le nouveau standard de traitement (NOACs)

EQUIPE MULTIDISCIPLINAIRE

First Amplatz in
Canada (2009)

First Watchman
in Canada (2012)

Merci!